arXiv:1412.3621v3 [math.OA] 27 Aug 2015

ON C*-ALGEBRAS WHICH CANNOT BE DECOMPOSED INTO TENSOR PRODUCTS WITH BOTH FACTORS INFINITE-DIMENSIONAL

TOMASZ KANIA

Abstract. We prove that C*-algebras which, as Banach spaces, are Grothendieck cannot be decomposed into a tensor product of two infinite-dimensional C*-algebras. By a result of Pfitzner, this class contains all von Neumann algebras and their norm-quotients. We thus complement a recent result of Ghasemi who established a similar conclusion for the class of SAW*-algebras.

1. Introduction and statement of the main result

During the London Mathematical Society Meeting held in Nottingham on 6$^{\text{th}}$ September 2010, Simon Wassermann asked a question of whether the Calkin algebra can be decomposed into a C*-tensor product of two infinite-dimensional C*-algebras. This question stems from the study of the elusive nature of the automorphism group of the Calkin algebra whose structure is independent of the usual axioms of Set Theory ([7, 18]). Ghasemi ([9]) studied tensorial decompositions of SAW*-algebras answering the above-mentioned question in the negative—one cannot thus expect to build automorphisms of such algebras out of automorphisms of non-trivial tensorial factors. Let us remark that the commutative version of Ghasemi's result was known to experts as it follows directly from the conjunction of [24, Theorem B] with the main theorem of [4].

The aim of this note is to prove that C*-algebras which satisfy a certain Banach-space property cannot be decomposed into a tensor product of C*-algebras. More specifically, we prove that C*-algebras which, as Banach spaces, are Grothendieck (*i.e.*, weak*-null sequences in the dual space converge weakly) do not allow such a tensorial decomposition. In particular, we give a new solution to the problem of Wassermann as the Calkin algebra falls into the class of Grothendieck spaces.

Theorem 1.1. *Let A be a C*-algebra which, as a Banach space, is a Grothendieck space. Suppose that E and F are C*-algebras such that*

$$A \cong E \otimes_\gamma F$$

for some C-norm γ. Then either E or F (or both) are finite-dimensional.*

In other words, a C-algebra, which is also a Grothendieck space, cannot be decomposed into a tensor product of two infinite-dimensional C*-algebras.*

Let us list examples of classes of C*-algebras which meet the assumptions of Theorem 1.1.

Proposition 1.2. *C*-algebras in each of the following classes are Grothendieck spaces:*

(i) *von Neumann algebras (or more generally, AW*-algebras) and their norm-quotients; in particular $\mathscr{B}(H)$ and the Calkin algebra,*
(ii) *ultraproducts of C*-algebras over countably incomplete ultrafilters,*
(iii) *unital C*-algebras with the countable Riesz interpolation property,*

Proof. By Corollary 2.4, von Neumann algebras and hence their continuous, linear images (such as the Calkin algebra) satisfy the hypothesis of Theorem 1.1. The assertion for AW*-algebras follows from Proposition 2.5 as every maximal abelian self-adjoint subalgebra of an AW*-algebra is Grothendieck by the main results of [22] and [23].

Avilés *et al.* ([2, Proposition 3.3]) proved that ultraproducts of Banach spaces over countably incomplete ultrafilters cannot contain complemented copies of c_0. This, combined with Theorem 2.2 and Proposition 2.1(iii), yields that ultraproducts of C*-algebras are Grothendieck spaces.

The assertion (iii) follows from applying [19, Theorem 9] to the real Banach space A_{sa} of all self-adjoint elements of a unital C*-algebra A with the countable Riesz interpolation property and noticing that the Grothendieck property passes from A_{sa} to the complex Banach space $A = A_{\mathrm{sa}} \oplus iA_{\mathrm{sa}}$. □

It is perhaps worthwhile to mention that even at the abelian level there exist many Grothendieck $C(X)$-spaces that are not SAW* (*i.e.*, for which X is not sub-Stonean); an example of such is a space constructed by Haydon ([12]).

Each unital C*-algebra with the countable Riesz interpolation property is an SAW*-algebra in the sense of Pedersen ([25, Proposition 2.7]); see [15] for the definition of an SAW*-algebra. Conjecturally all unital SAW*-algebras have the countable Riesz interpolation property ([25, p. 117]), hence our result covers all known examples of SAW*-algebras—thus, we extend the main result of [9] to the class of ultraproducts of C*-algebras and other C*-algebras created, for instance, out of Grothendieck abelian C*-algebras that are not SAW*.

To the best of our knowledge, it is not known whether a maximal abelian self-adjoint subalgebra of an SAW*-algebra is SAW* too. If this were the case, Proposition 2.5 would immediately imply that SAW*-algebras are Grothendieck spaces, because abelian SAW*-algebras are of the form $C_0(X)$ for some locally compact sub-Stonean space X, hence Grothendieck by [1] or [24].

2. Preliminaries

Grothendieck spaces. A series $\sum_{n=1}^{\infty} y_n$ in a Banach space E is *weakly unconditionally convergent* if the scalar series $\sum_{n=1}^{\infty} |\langle f, y_n\rangle|$ converges for each $f \in E^*$. An operator between Banach spaces is *unconditionally converging* if it maps weakly unconditionally convergent series to unconditionally convergent series. A Banach space E has *property (V)*

if for each Banach space F the class of unconditionally converging operators $T\colon E \to F$ coincides with the class of weakly compact operators. It is a result of Pełczyński that $C(X)$-spaces (abelian C*-algebras) have property (V) ([16]).

A Banach space E is *Grothendieck* if every weak*-null sequence in E^* converges weakly. The name *Grothendieck space* stems from a result of Grothendieck ([11]) who identified ℓ_∞ as a space having this property. It is a trivial remark that reflexive spaces have this property too. By the Hahn–Banach theorem, the class of Grothendieck spaces is closed under surjective linear images, *i.e.*, whenever E and F are Banach spaces, $T\colon E \to F$ is a surjective bounded linear operator, then if E is Grothendieck, so is F.

Let us record a proposition which links property (V) with the Grothendieck property.

Proposition 2.1. *Let X be a Banach space. Then the following are equivalent.*

(i) *X is a Grothendieck space,*
(ii) *each bounded linear operator $T\colon X \to c_0$ is weakly compact.*
(iii) *X has property (V) and no subspace of X isomorphic to c_0 is complemented.*

Proof. For the proof of equivalences (i) $\iff$ (ii) see [5, Corollary 5 on p. 150]. Equivalence (i) $\iff$ (iii) is due to Räbiger ([20]); see also [10, Theorem 28] (this argument is also implicit in the proof of [4, Corollary 2]). □

We require the following theorem of Pfitzner ([17, Theorem 1]), which can be thought as a non-commutative generalisation of the above-mentioned result of Pełczyński.

Theorem 2.2 (Pfitzner)**.** *Let A be a C*-algebra and let $\mathcal{K} \subset A^*$ be a bounded set. Then $\mathcal{K}$ is not relatively weakly compact if and only if there are a sequence $(x_n)_{n=1}^\infty$ of pairwise orthogonal, norm-one self-adjoint elements in A and $\delta > 0$ such that*

$$\sup_{f\in\mathcal{K}} |\langle f, x_n\rangle| > \delta.$$

In particular, C-algebras have property (V).*

The original proof was highly sophisticated and relied on numerous deep facts from Banach space theory. Fortunately, Fernández-Polo and Peralta ([8]) supplied a short and elementary proof of Pfitzner's theorem.

By virtue of Proposition 2.1(iii), we arrive at the following corollary.

Corollary 2.3. *A C*-algebra is a Grothendieck space if and only if it does not contain complemented subspaces isomorphic to c_0.*

The special case of Corollary 2.3 where the C*-algebra is also a von Neumann algebra was noted by Pfitzner ([17, Corollary 7]):

Corollary 2.4. *Von Neumann algebras are Grothendieck spaces.*

Let us take this opportunity to record the following easy corollary to Theorem 2.2.

Proposition 2.5. *Let A be a C*-algebra with the property that each maximal abelian self-adjoint subalgebra B of A is a Grothendieck space. Then A is a Grothendieck space.*

Proof. Let $T\colon A \to c_0$ be a bounded linear operator. By Proposition 2.1(ii) it is enough to show that T is weakly compact.

Assume contrapositively that T is not weakly compact. By Gantmacher's theorem, T is weakly compact if and only if T^* is, so the set $\mathcal{K} = T^*(B)$ is not relatively weakly compact, where B is the unit ball of c_0^*. By Theorem 2.2, there exist $\delta > 0$ and a sequence $(x_n)_{n=1}^{\infty}$ of pairwise orthogonal, norm-one self adjoint elements in A such that

$$\sup_{f\in\mathcal{K}} |\langle f, x_n\rangle| = \sup_{y\in B} |\langle T^*y, x_n\rangle| = \sup_{y\in B} |\langle y, Tx_n\rangle| > \delta. \tag{2.1}$$

Let $B_0 \subseteq A$ be the C*-algebra generated by $\{x_n\colon n \in \mathbb{N}\}$. Since the x_n ($n \in \mathbb{N}$) are pairwise orthogonal, B_0 is abelian. Let B be a maximal abelian subalgebra of A containing B_0. Consider the restriction $T|_B\colon B \to c_0$. It is not weakly compact by (2.1), so B is not a Grothendieck space. □

3. Proof of Theorem 1.1

We are now in a position to prove our main result. The general strategy of the proof was inspired by the path taken by Cembranos in [4].

Proof of Theorem 1.1. Let A be a C*-algebra and suppose that it is a Grothendieck space. Assume towards a contradiction that $A \cong E\otimes_\gamma F$ for some infinite-dimensional C*-algebras E, F and a C*-norm γ.

Since *-homomorphisms between C*-algebras have closed range, there is a surjective *-homomorphism $Q\colon E\otimes_\gamma F \to E\otimes_{\min} F$ that extends the identity map on $E\odot F$. (Here $E\otimes_{\min} F$ denotes the minimal C*-tensor product of E and F.)

Let $B_1 \subset E$ and $B_2 \subset F$ be infinite-dimensional abelian C*-algebras. (Such subalgebras exist because infinite-dimensional C*-algebras contain self-adjoint elements with infinite spectrum ([13, Ex. 4.6.12]), hence the assertion follows from the spectral theorem.) We may thus identify $B_1\otimes_{\min} B_2$ with a subalgebra of $E\otimes_{\min} F$ (*cf.* [3, II.9.6.2]). However, the (minimal) tensor product of abelian C*-algebras is the same as the Banach-space injective tensor product, *i.e.*,

$$B_1 \otimes_{\min} B_2 = B_1 \mathbin{\check{\otimes}} B_2.$$

Let $(e_n)_{n=1}^{\infty}$ be a sequence of pairwise orthogonal, positive, norm-one elements in B_1. In particular, $E_0 = \overline{\operatorname{span}}\{e_n\colon n \in \mathbb{N}\}$ is isometric to c_0 and $(e_n)_{n=1}^{\infty}$ is equivalent to the canonical basis for c_0. Choose norm-one functionals $e_n^* \in E^*$ such that $\langle e_n^*, e_m\rangle = \delta_{n,m}$ ($n, m \in \mathbb{N}$). Moreover, let $(x_n^*)_{n=1}^{\infty}$ be a sequence of unit vectors in F^* which converges to 0 in the weak* topology. (Such a sequence exists by the Josefson–Nissenzweig theorem (see [6, Chapter XII].) Let $(x_n)_{n=1}^{\infty} \subset F$ be a sequence such that $\langle x_n, x_n^*\rangle = 1$. Without loss of generality we may suppose that $\|x_n\| \leqslant 2$ for all n.

Define a map $T\colon E\odot_{\min} F \to \ell_\infty$ by the formula

$$T\xi = (\langle e_n^*\otimes x_n^*, \xi\rangle)_{n=1}^{\infty} \qquad (\xi \in E\odot_{\min} F).$$

This is a well-defined bounded linear operator because $(e_n^* \otimes x_n^*)_{n=1}^\infty$ is a bounded sequence of functionals on $E \odot_{\min} F$. We can thus extend T to the whole of $E \otimes_{\min} F$. Moreover, for all $f \in E$ and $x \in F$ we have

$$|\langle e_n^*, f\rangle \cdot \langle x, x_n^*\rangle| \leqslant \|f\| \cdot |\langle x, x_n^*\rangle|$$

so T takes values in c_0 as $(x_n^*)_{n=1}^\infty$ is a weak*-null sequence. Since the injective tensor product 'respects subspaces' (see [21, p. 49]), $E_0 \mathbin{\check{\otimes}} B_2$ can be identified with a subspace of $B_1 \mathbin{\check{\otimes}} B_2$ and the latter is a subspace of $E \otimes_{\min} F$.

As E_0 and c_0 are isometrically isomorphic, so are $E_0 \mathbin{\check{\otimes}} B_2$ and $c_0 \mathbin{\check{\otimes}} B_2 \cong c_0(B_2)$ (*cf.* [21, Example 3.3]). Let $n \in \mathbb{N}$ and let $(a_k)_{k=1}^\infty$ be a scalar sequence with only finitely many non-zero entries. We have

$$\Big\| \sum_{k=1}^n a_k e_k \otimes x_k \Big\| = \Big\| \sum_{k=1}^n e_k \otimes (a_k x_k) \Big\| \leqslant \sup_{1 \leqslant k \leqslant n} \|a_k x_k\| \leqslant 2 \max\{|a_k| \colon 1 \leqslant k \leqslant n\}.$$

By [14, Proposition 4.3.9], $\sum_{n=1}^\infty e_n \otimes x_n$ is a weakly unconditionally convergent series in $E \otimes_{\min} F$. On the other hand, for all $k, n \in \mathbb{N}$ we have $(T(e_n \otimes x_n))(k) = \delta_{k,n}$ so

$$\sum_{n=1}^\infty T(e_n \otimes x_n)$$

fails to converge in c_0. Consequently, $E \otimes_{\min} F$ is not a Grothendieck space as we proved that the c_0-valued operator T is not unconditionally converging. Indeed, if $E \otimes_{\min} F$ were Grothendieck, T would be weakly compact (Proposition 2.1(ii)), hence also unconditionally converging (Proposition 2.1(iii)). □

Acknowledgments. We are indebted to Y. Choi and N. Ozawa for spotting an error in the previous version of the proof of Theorem 1.1.

References

1. T. Andô, Convergent sequences of finitely additive measures, *Pacific J. Math.* **11** (1961), 395–404.
2. A. Avilés, F. Cabello Sánchez, J. M. F. Castillo, M. González and Y. Moreno, On ultrapowers of Banach spaces of type $\mathscr{L}_\infty$, *Fund. Math.* **222** (2013), 195–212.
3. B. Blackadar, *Operator algebras*, Encyclopaedia of Mathematical Sciences, vol. 122, Springer-Verlag, Berlin, 2006, Theory of C*-algebras and von Neumann algebras, Operator Algebras and Non-commutative Geometry, III.
4. P. Cembranos, $C(K, E)$ contains a complemented copy of c_0, *Proc. Amer. Math. Soc.* **91** (1984), 556–558.
5. J. Diestel, *Geometry of Banach Spaces*. Selected Topics, Lecture Notes in Math. 485, Springer (1975).
6. J. Diestel, *Sequences and Series in Banach Spaces*, Volume 92 of Graduate Text in Mathematics. Springer Verlag, New York, First edition, (1984).
7. I. Farah, All automorphisms of the Calkin algebra are inner, *Annals of Mathematics*, **173** (2010), 619–661.
8. F.J. Fernández-Polo and A.M. Peralta, A short proof of a theorem of Pfitzner, *Q. J. Math.* **61** (2010), 329–336.
9. S. Ghasemi, SAW*-algebras are essentially non-factorizable, *Glasgow Math. J.* **57** (2015), 1–5.
10. I. Ghenciu and P. Lewis, Completely continuous operators, *Colloq. Math.* **126**, No. 2 (2012), 231–256.

11. A. Grothendieck, Sur les applications linéaires faiblement compactes d'espaces du type $C(K)$, *Canadian J. Math.* **5** (1953), 129–173.
12. R. Haydon, A non-reflexive Grothendieck space that does not contain ℓ_∞, *Israel J. Math.*, **40** (1981), 65–73.
13. R. V. Kadison and J. R. Ringrose, *Fundamentals of the Theory of Operator Algebras, Vol. I, Elementary Theory*, Pure and Applied Math., Vol. 100 Academic Press, New York (1983).
14. R. E. Megginson, *An Introduction to Banach Space Theory*, Graduate Texts in Mathematics 183, Springer-Verlag, 1998.
15. G. K. Pedersen, SAW*-algebras and corona C*-algebras, contributions to non-commutative topology, *J. Operator Theory*, **15** (1986), 15–32.
16. A. Pełczyński, On Banach spaces on which every unconditionally converging operator is weakly compact, *Bull. Acad. Pol. Sci.*, **10** (1962), 641–648.
17. H. Pfitzner, Weak compactness in the dual of a C*-algebra is determined commutatively, *Math. Ann.*, **298** (1994), 349–371.
18. N.C. Phillips and N. Weaver, The Calkin algebra has outer automorphisms, *Duke Math. J.* **139** (2007), 185–202.
19. I. A. Polyrakis and F. Xanthos, Grothendieck ordered Banach spaces with an interpolation property, *Proc. Amer. Math. Soc.* **141** (2013), 1651–1661.
20. F. Räbiger, *Beiträge zur Strukturtheorie der Grothendieck-Räume*, Sitzungsber., Heidelberger Akad. Wiss., Math.-Naturwiss. Kl. 4, 78 S.
21. R. A. Ryan, *Introduction to Tensor Products of Banach Spaces*, Springer, 2002.
22. K. Saitô and J.D. Maitland Wright, C^*-algebras which are Grothendieck spaces, *Rend. Circ. Mat. Palermo* **52** (2003), 141–144.
23. K. Saitô and J.D. Maitland Wright, On Defining AW*-algebras and Rickart C*-algebras, to appear in *Q. J. Math*, doi:10.1093/qmath/hav015.
24. G. L. Seever, Measures on F-spaces, *Trans. Amer. Math. Soc.*, **133** (1968), 267–280.
25. R. Smith and D. Williams, Separable injectivity for C*-algebras, *Indiana Univ. Math. J.*, **37** (1988) 111–133.
26. S. Wassermann, Talk at the London Mathematical Society Meeting held in Nottingham on 6$^{\text{th}}$ September 2010.

Department of Mathematics and Statistics, Fylde College, Lancaster University, Lancaster LA1 4YF, United Kingdom.

E-mail address: tomasz.marcin.kania@gmail.com